

Journal of Drug Discovery and Therapeutics

Available Online at www.jddt.in

CODEN: - JDDTBP (Source: - American Chemical Society)

Volume 11, Issue 06: 2023, 56-63

Medicinal Plants Used for Diabetes Mellitus: A Short Review

Prashant Kumar^{1*} & Richa Mishra²

¹ Research Scholar, Department of Pharmacy, Sunrise University, Alwar Rajasthan.

² Research Supervisor, Department of Pharmacy, Sunrise University, Alwar Rajasthan.

Received: 07-08-2023 / Revised: 14-09-2023 / Accepted: 09-10-2023

Corresponding author: Prashant Kumar

Conflict of interest: No conflict of interest.

Abstract:

Investigating the antidiabetic effects of different medicinal plants is the focus of the current study. Among the many non-communicable illnesses that affect people across the world, diabetes mellitus ranks high. Although it has not yet reached pandemic proportions in many poor and newly industrialized nations, it is now the fourth worst killer among the world's wealthiest countries. We must confront this grave danger in the twenty-first century. Herbs and plants have been used as medicine for a very long time. Plants were referenced in Ayurveda and other Indian texts as potential treatments for a variety of illnesses. Less than one percent of the approximately 250,000 higher plants have undergone pharmacological screening, with even less tested for diabetes mellitus. There has been a dearth of systematic research on the traditional medicinal herbs used to treat diabetes mellitus.

Introduction

Developed and developing world residents alike are not immune to the effects of diabetes mellitus. This illness is thought to impact over 25% of the global population. When insulin levels in the blood are low or when insulin is unable to reach its target organs, a metabolic disorder known as diabetes mellitus develops [1]. The current synthetic medications have a number of limitations, thus the hunt for additional drugs continues, even if oral hypoglycemic agents have made great strides in the treatment of diabetes. Despite their widespread praise for their medicinal efficacy in traditional medicine, herbal medications with antidiabetic action have not yet been developed for commercial use as contemporary pharmaceuticals [2].

Obesity is a risk factor for type 2 diabetes, which in turn increases the risk of hypertension and dyslipidemia. Therefore, the goal of the therapy is to increase insulin secretion and decrease insulin resistance. A metabolic illness known as diabetes occurs when the body either does not create enough insulin or does not use it correctly. Insulin is a hormone that the body needs to convert food sources like sugar and carbs into energy. In diabetes mellitus, blood glucose (sugar) levels are consistently high. Glucagon and insulin work together to keep blood glucose levels in the human body within a tight range. Glucagon stimulates the release of glucose from the liver's cells into the bloodstream, which is essential for energy generation. Muscles and adipose

tissue absorb glucose at low rates due to Type 1 Diabetes, which is characterised by an inability to secrete insulin [3]. Because of the high expense of contemporary medications, people in underdeveloped nations often turn to traditional medicine, namely herbal remedies, to manage their diabetes [4]. Diabetes and its consequences remain a significant health concern, even with the development of hypoglycemic drugs derived from both natural and synthetic sources. Diabetes may be effectively managed with the use of many native Indian medicinal herbs. The minimal risk of adverse effects and wide availability of medicinal plants are two of their greatest strengths. Numerous pharmaceuticals that are accessible today have their origins in plants, and this trend has persisted throughout history.

Approximately 800 plants have been identified in ethnobotanical studies as

having antidiabetic properties [5]. Current experimental methods have shown antidiabetic efficacy in a number of botanicals [6]. This review article enumerates some medicinal plants possessing antidiabetic activity and elucidating their mechanisms of action such as *Brassica juncea* (*B. juncea*), *Combretum micranthum* (*C. micranthum*), *Elephantopus scaber* (*E. scaber*), *Gymnema sylvestre* (*G. sylvestre*), *Liriope spicata* (*L. spicata*), *Parinari excelsa* (*P. excelsa*), *Ricinus communis* (*R. communis*), *Sarcopoterium spinosum* (*S. spinosum*), *Smallanthus sonchifolius* (*S. sonchifolius*), *Swertia punicea* (*S. punicea*), *Vernonia anthelmintica* (*V. anthelmintica*) etc. and method of experiment on animals and therapeutic efficiency of plant extracts were exploited. Table 1 lists some of the most promising herbal plant sources for anti-diabetic effects.

Table 1: Medicinal plants having antidiabetic activity.

| Plantname | Family | Parts used | Typeofextract | Activity | References |
|---------------------------------|----------------|-------------|----------------------|-----------------------------------|------------|
| <i>Alangium lamarckii</i> | Alangiaceae | Leaves | Alcoholic | Antidiabetic | [15] |
| <i>Albizia odoratissima</i> | Mimosaceae | Bark | Methanol | Antidiabetic | [19] |
| <i>Axonopus compressus</i> | Poaceae | Leaves | Methanol | Antidiabetic | [13] |
| <i>Berberis vulgaris</i> | Berberidaceae | Root | Aqueous | Hypoglycaemic | [14] |
| <i>Brassica juncea</i> | Cruciferae | Seed | Aqueous | Hypoglycemic | [7] |
| <i>Caesalpinia digyna</i> | Fabaceae | Root | Methanol | Antidiabetic | [10] |
| <i>Catharanthus roseus</i> | Apocynaceae | Leaf | Methanol | Hypoglycemic | [16] |
| <i>Centaurium erythraea</i> | Gentianaceae | Leaf | Aqueous | Antidiabetic | [17] |
| <i>Chaenomeles sinensis</i> | Rosaceae | Fruits | ethylacetate | Antidiabetic | [18] |
| <i>Cocos nucifera</i> | Arecaceae | Leaf | hydro-methanol | Antihyperglycemic | [20] |
| <i>Costus speciosus</i> | Costaceae | rhizome | hexane | Antidiabetic | [21] |
| <i>Cyclocarya paliurus</i> | Cyclocaryaceae | Bark | Aqueous | Hypoglycemic | [22] |
| <i>Dillenia indica</i> | Dilleniaceae | Leaves | Methanolic | Antidiabetic | [23] |
| <i>Embelia ribes</i> | Myrsinaceae | Berries | Hexane | Antidiabetic | [24] |
| <i>Hybanthus enneaspermus</i> | Violaceae | Whole plant | Alcoholic | Antidiabetic | [25] |
| <i>Lippa nodiflora</i> | Verbenaceae | Whole plant | Methanol | Antidiabetic Hypolipidemic | [26] |
| <i>Lithocarpus polystachyus</i> | Fagaceae | Leaves | Ethanol & Aqueous | Hypoglycemic | [27] |
| <i>Marrubium vulgare</i> | Lamiaceae | Aerial part | Methanol | Hyperglycemia and dyslipidemia | [28] |

| | | | | | |
|----------------------------------|-----------------|--------------|----------------------|-------------------|------|
| <i>Ocimum sanctum</i> | Lamiaceae | Aerial part | Hydroalcoholic | Antidiabetic | [29] |
| <i>Opuntia streptacantha</i> | Cactaceae | Leaves | Ethanol | Antihyperglycemia | [30] |
| <i>Psidium guajava</i> | Myrtaceae | Fruits | Ethanol | Antihyperglycemic | [31] |
| <i>Semecarpus anacardium</i> | Anacardiaceae | nut | Milk | Antidiabetic | [32] |
| <i>Prosopis glandulosa</i> | Fabaceae | Whole plant | Gelatine/Jelly | Antidiabetic | [33] |
| <i>Ophiopogon japonicus</i> | Asparagaceae | Root | Ethanol | Hypoglycemic | [34] |
| <i>Setaria italica</i> | Poaceae | Seed | Aqueous | Antihyperglycemic | [35] |
| <i>Solanum torvum</i> | Solanaceae | Fruit | Methanol | Antidiabetic | [36] |
| <i>Cassia auriculata</i> | Caesalpiniaceae | Leaves | Aqueous | Antidiabetic | [37] |
| <i>Zygodium album</i> | Zygophyllaceae | Whole plant | Ethanol | Antihyperglycemic | [38] |
| <i>Vitex negundo</i> | Lamiaceae | Leaves | Methanol | Antidiabetic | [39] |
| <i>Viscum schimperi</i> | Viscaceae | aerial parts | Methanolic | Antidiabetic | [40] |
| <i>Symplocos cochinchinensis</i> | Symplocaceae | Leaves | Hexane | Antihyperglycemic | [41] |
| <i>Enicostemma littorale</i> | Gentianaceae | Whole plant | aqueous | Antidiabetic | [42] |
| <i>Vaccinium arctostaphylos</i> | Ericaceae | Fruit | Ethanol | Antidiabetic | [43] |
| <i>Solanum xanthocarpum</i> | Solanaceae | Leaves | Aqueous and Methanol | Antidiabetic | [44] |

According to traditional medicine, *E. jambolana* has anti-diabetic benefits. In both experimental models and clinical trials, it has been shown to have hypoglycemic effects [8].

Antidiabetic effect of folklore medicinal plants

Brassica juncea

As a spices, it finds widespread use in Tamilnadu cuisine. The traditional medicinal herb *B. juncea* is a member of the Cruciferae family. Researchers looked at the hypoglycemic effects of an aqueous seed extract from *B. juncea* in male albino rats that had been induced diabetes by streptozotocin. There have been reports of doses of 250, 350, and 450 mg/kg having hypoglycemic action [7].

Coccinia grandis

Coccinia grandis (*C. grandis*) leaf alcoholic extracts were tested for hypoglycemic activity. Mice were gavaged with an alcoholic extract at a dose of 600 mg/kg body weight. In normal, fasting mice, an alcoholic extract of *C. grandis* leaves given orally significantly reduced blood glucose levels [9].

Alangium lamarckii

Antidiabetic action of alcoholic extract of *Alangium lamarckii* (*A. lamarckii*). Both the 250 mg/kg bw and 500 mg/kg bw investigations used alcoholic leaf extract. The antidiabetic effects of *A. lamarckii* were seen in rats who were diabetic after being administered STZ-nicotinamide[10].

Albizia odoratissima

Methanolic bark extract of *Albizia odoratissima* (*A. odoratissima*) and its potential anti-diabetic effects in diabetic rats produced by alloxan has been studied. The dosages given to the animals for the methanolic extracts were 250 and 500 mg/kg body weight. In albino mice induced with alloxan, serum cholesterol, triglycerides, SGOT, SGPT, alkaline phosphatase, and total proteins were significantly decreased [11].

Artemis sphaerocephala Krasch

Artemis sphaerocephala (*A. sphaerocephala*) gum's antioxidant effects in diabetic rats

produced by streptozotocin (STZ). The levels of +OH and thiobarbituric acid reactive substances (TBARS) in the serum and liver of STZ-induced rats were upregulated. Superoxide dismutase activity in the liver and blood was reduced. There was a reduction in blood and liver tissue levels of TBARS and +OH after treatment of *A. sphaerocephala* extract. The substantial increases in superoxide dismutase (SOD) levels in the liver and blood. The antioxidant activity of *A. sphaerocephala* is excellent [12].

Axonopus compressus

How the plant's methanolic leaf extract prevents diabetes. Injecting the rats with alloxan caused them to develop diabetes. The experiments used methanolic leaves extract at concentrations of 250, 500, and 1,000 mg/kg bw. When compared to the control group, diabetic rats treated with methanolic leaf extract of *Axonopus compressus* (*A. compressus*) at dosages of 250, 500, and 1,000 mg/kg showed a substantial decrease (31.5%, 19.8%, and 24.5%, respectively) in blood glucose levels. It is possible that *A. compressus* has excellent antidiabetic properties [13].

Berberis vulgaris

Herb *Berberis vulgaris* (*B. vulgaris*) L.'s hypoglycemic effects in diabetic rats produced by streptozotocin. The Berberidaceae family includes the traditionally used medicinal herb *B. vulgaris*. The findings demonstrated a significant hypoglycemic impact of the water extract and saponins. Serum triglyceride and cholesterol levels were much higher than normal [14].

Caesalpinia digyna

Bergenin, derived from *Caesalpinia digyna* (*C. digyna*) roots, has an antidiabetic action. Diabetic rats had much higher plasma total cholesterol (TC), triglycerides (TG), and

low-density lipoprotein (LDL-C) levels than control rats, but much lower levels of good cholesterol (HDL-C). The lipid profile was noticeably altered after glibenclamide (10 mg/kg; p.o.) treatment compared to bergenin (10 mg/kg; p.o.). Antioxidant enzyme activity, including SOD and Cat, was reduced. Diabetic rats have far higher levels of TBARS than control rats. Increasing SOD and CAT levels while decreasing TBARS was a notable effect of bergenin (10 mg/kg; p.o.). There are excellent antidiabetic activities in bergenin [15].

Catharanthus roseus

Investigation into the hypoglycemic effects of *Catharanthus roseus* (*C. roseus*) methanolic leaf extract in rats induced with diabetes by alloxan. Blood glucose levels were much lower than those of the control group of rats. *C. roseus* methanolic extract had a stronger impact in lowering blood glucose levels compared to Glibenclamide and Metformin [16].

Centaurium erythraea

A single intraperitoneal injection of 65 mg/kg of STZ caused diabetes. A measure of oxidative stress was performed using tissue MDA. The assessment of antioxidant enzymes produced by the pancreas, including SOD, CAT, and GPx (glucose peroxidase). Pancreatic tissue TBARS levels were significantly lower in diabetes treated rats compared to control animals. Diabetic animals treated with antioxidant defence enzymes showed a substantial rise in activity levels of SOD, CAT, GPx, and GST. *Centaurium erythraea* (*C. erythraea*) aqueous leaf extract's antioxidant activity [17].

Chaenomeles sinensis

An excellent antidiabetic effect is produced by the ethyl acetate fraction of the fruits of *Chaenomeles sinensis* (*C. sinensis*) (Thouin) Koehne. The Rosaceae family includes *Chaenomeles sinensis*. Doses of 50 and 100

mg/kg body weight were shown to have antidiabetic effects [18].

Conclusion

Traditional medicinal plants used in folk medicine to treat diabetes mellitus were included in this study. Traditional medicinal herbs are more often employed in rural regions due to the abundance of therapeutic plants found there. As a result, the prospect of treating diabetes mellitus using chemicals derived from plants that are both accessible and do not need the time-consuming pharmaceutical production is quite appealing. Health care providers, researchers, and academics in the domains of pharmacology and therapeutics may find this review helpful as they endeavour to learn more about antidiabetic medicinal plants.

References

1. Ajay SS. Hypoglycemic activity of *Coccinia indica* (Cucurbitaceae) leaves. *Int J Pharm Tech Res* 2009; 1(3): 892-893.
2. AL Lehninger, DL Nelson, MM Cox. *Principle of Biochemistry*. New York: Worth Publishers; 2010.
3. Alarcon-Aguilara FJ, Roman-Ramos R, Perez-Gutierrez S, Aguilar-Contreras A, Contreras-Weber CC, Flores-Saenz JL. Study of the anti-hyperglycemic effect of plants used as antidiabetics. *J Ethnopharmacol* 1998; 61: 101-110.
4. Balamurugan R, Ignacimuthu S. Antidiabetic and hypolipidemic effect of methanol extract of *Lippia nodiflora* L. in STZ induced diabetic rats. *Asian Pac J Trop Biomed* 2011; 1: S30-36.
5. Cetto AA, Wiedenfeld H. Anti-hyperglycemic effect of *Opuntia streptacantha* Lem. *J Ethnopharmacol* 2011; 133: 940-943.
6. Chen X, Jin J, Tang J, Wang Z, Wanga J, Jin L, et al. Extraction, purification, characterization and hypoglycemic activity of a polysaccharide isolated from the root of *Ophiopogon japonicas*. *Carbohydrate Polymers* 2011; 83: 749-754.
7. Dinesh Kumar, Sunil Kumar, Sonia Kohli, Renu Arya, Jyoti Gupta. Antidiabetic activity of methanolic bark extract of *Albizia odoratissima* Benth. in alloxan induced diabetic albino mice. *Asian Pac J Trop Med* 2011; 900-903.
8. Elberry AA, Harraz FM, Ghareib SA, Gabr SA, Nagy AA, Sattar EA. Methanolic extract of *Marrubium vulgare* ameliorates hyperglycemia and dyslipidemia in streptozotocin-induced diabetic rats. *Int J Diabetes Mellitus* (2011). In press.
9. Eliza J, Diasy P, Ignacimuthu S, Duraipandiyani V. Antidiabetic and antilipidemic effect of eremanthin from *Costus speciosus* (Koen.)Sm., in STZ-induced diabetic rats. *Chem Biol Interact* 2009; 182: 67-72.
10. Feshani AM, Kouhsari SM, Mohammadi S. *Vaccinium arctostaphylos*, a common herbal medicine in Iran: Molecular and biochemical study of its antidiabetic effects on alloxan- diabetic Wistar rats. *J Ethnopharmacol* 2011; 133: 67-74.
11. Gandhi GR, Ignacimuthu S, Paulraj MG, Sasikumar P. Antihyperglycemic activity and antidiabetic effect of methyl caffeate isolated from *Solanum torvum* Swartz. fruit in streptozotocin induced diabetic rats. *Eur J Pharmacol* 2011; 670: 623-631.
12. Georgea C, Lochnera A, Huisamen B. The efficacy of *Prosopis glandulosa* as antidiabetic treatment in rat models of diabetes and insulin resistance. *J Ethnopharmacol* 2011; 137: 298-304.
13. Ghoul JE, Boughanmi NG, Attia MB. Biochemical study on the protective effect of ethanolic extract of *Zygophyllum album* on streptozotocin induced oxidative stress and toxicity in

- mice. *Biomed Preventive Nutr* 2011; 1(2): 79-83.
14. Gupta S, Sharma SB, Singh UR, Bansal SK. Salutory effect of *Cassia auriculata* L. leaves on hyperglycemia-induced atherosclerotic environment in streptozotocin rats. *Cardiovasc Toxicol* 2011; 11: 308-315.
 15. Hedayathullah Khan HB, Vinayagam KS, Palanivelu S, Panchanatham S. Anti-diabetic effect of *Semecarpus anacardium* Linn nut milk extract in a high fat diet STZ- induced type 2 diabetic rat model. *Comp Clin Pathol* 2012; 21(6): 1395-1400.
 16. Hou SZ, Chen SX, Huang S, Jiang DX, Zhou CJ, Chen CQ, et al. The hypoglycemic activity of *Lithocarpus polystachyus* Rehd. leaves in the experimental hyperglycemic rats. *J Ethnopharmacol* 2011; 138: 142- 149.
 17. Huang CS, Yin MC, Chiu LC. Antihyperglycemic and antioxidative potential of *Psidium guajava* fruit in streptozotocin-induced diabetic rats. *Food Chem Toxicol* 2011; 41: 2189-2195.
 18. Ibeh BO, Ezeaja MI. Preliminary study of antidiabetic activivty of the methanolic leaf extract of *Axonopus compressus* (P.Beauv) in alloxan induced diabetic rats. *J Ethnopharmacol* 2011; 138: 713-716.
 19. Jafri MA, Aslam M, Javed K, Singh S. Effect of *Punica granatum* Linn (flowers) on blood glucose level in normal and alloxan-induced diabetic rats. *J Ethnopharmacol* 2000; 70:309-314.
 20. Kumar D, Kumar S, kohli S, Arya R, Gupta J. Antidaibetic activity of methanolic bark extract of *Albizia odoratissima* Benth in alloxan induced diabetic albino mice. *Asian Pac J Trop Med* 2011; 4: 900-903.
 21. Kumar R, Patel DK, Prasad SK, Laloo D, Krishnamurthy S, Hemalatha S. Type 2 antidiabetic activity of bergenin from the roots of *Caesalpinia digyna* Rottler. *Fitoterapia* 2012; 83(2): 395-401.
 22. Kumar S, Kumar V, Om Prakash. Antidiabetic, hypolipidemic and histopathological analysis of *Dillenia indica* (L.) leaves extract on alloxan induced diabetic rats. *Asian Pac J Trop Med* 2011; 347-352.
 23. Li S, Li J, Guan XL, Li J, Deng SP, Li LQ, et al. Hypoglycemic effects and constituents of the barks of *Cyclocarya paliurus* and their inhibiting activities to glucosidase and glycogen phosphorylase. *Fitoterapia* 2011; 82: 1081-1085.
 24. Mahendran S, Badami S, Maithili V. Evaluation of antidiabetic effect of embelin from *Embelia ribes* in alloxan induced diabetes in rats. *Biomed Preventive Nutr* 2011; 1: 25-31.
 25. Maiti R, Jana D, Das UK, Ghosh D. Antidiabetic effect of aqueous extract of seed of *Tamarindus indica* in streptozotocin induced diabetic rats. *J Ethnopharmacol* 2004; 92: 85-91.
 26. Meliani N, Amine Dib ME, Allali H, Tabti B. Hypoglycaemic effect of *Berberis vulgaris* L. in normal and streptozotocin induced diabetic rats. *Asian Pac J Trop Biomed* 2011; 6: 468-471.
 27. Naskar S, Mazumder UK, Pramanik G, Gupta M, Sureshkumar RB, Bala A, et al. Evaluation of antihyperglycemic activity of *Cocos nucifera* Linn.on streptozotocin induced type 2 diabetic rats. *J Ethnopharmacol* 2011; 138: 769-773.
 28. Ohadoma SC, Michael HU. Effects of co-administration of methanol leaf extract of *Catharanthus roseus* on the hypoglycemic activity of metformin and

- glibenclamide in rats. *Asian Pac J Trop Med* 2011; 475-477.
29. Patel DK, Kumar R, Prasad SK, Sairam K, Hemalatha S. Antidiabetic and in vitro antioxidant potential of *Hybanthus enneaspermus* (Linn) F. Muell in streptozotocin-induced diabetic rats. *Asian Pac J Trop Med* 2011; 4: 316-322.
 30. Patil R, Patil R, Ahirwar B, Ahirwar D. Isolation and characterization of anti-diabetic component (bioactivity guided fractionation) from *Ocimum sanctum* L. (Lamiaceae) aerial part. *Asian Pac J Trop Med* 2011; 278-282.
 31. Poongothai K, Ponnuragan P, Syed Zameer Ahmed K, Senthil Kumar B, Sheriff SA. Antihyperglycemic and antioxidant effects of *Solanum xanthocarpum* leaves (field grown & in vitro raised) extracts on alloxan induced diabetic rats. *Asian Pac J Trop Med* 2011; 778-785.
 32. Rajesh Kumar, Dinesh Kumar Pate, Satyendra Kuldip Prasad, Kirshnamurthy Sairam, Siva Hemalatha. Antidiabetic activity of alcoholic leaves extract of *Alangium lamarckii* Thwaites on streptozotocin-nicotinamide induced type 2 diabetic rats. *Asian Pac J Tropical Med* 2011; 904-909.
 33. Ravi K, Ramachandran B, Subramanian S. Effect of *Eugenia Jambolana* seed kernel on antioxidant defense system in streptozotocin induced diabetes in rats. *Life Sci* 2004; 75(22): 2717-2731.
 34. Sancheti S, Sancheti S, Seo SY. Antidiabetic and antiacetylcholinesterase effects of ethyl acetate fraction of *Chaenomeles sinensis* (Thouin) Koehne fruits in streptozotocin-induced diabetic rats. *Exp Toxicol Pathol* 2011; 65(1-2): 55-60.
 35. Saravanan G, Pari L. Hypoglycaemic and antihyperglycaemic effect of *Syzygium cumini* bark in streptozotocin-induced diabetic rats. *J Pharmacol Toxicol* 2008; 3: 1-10.
 36. Sattar EA, Elberry AA, Harraz FM, Ghareib SA, Nagy AA, Gabr SA. Antihyperglycemic and hypolipidaemic effects of the methanolic extract of Saudi mistletoe (*Viscum schimperi* Engl.). *J Adv Res* 2011; 2: 171-177.
 37. Sefi M, Fetoui H, Lachkar N, Tahraoui A, Lyoussi B, Boudawara T, et al. *Centaurium erythraea* (Gentianaceae) leaf extract alleviates streptozotocin-induced oxidative stress and β -cell damage in rat pancreas. *J Ethnopharmacol* 2011; 135: 243-250.
 38. Sireesh Y, Kasetti RB, Nabi SA, Swapna S, Apparao C. Antihyperglycemic and hypolipidemic activities of *Setaria italica* seeds in STZ diabetic rats. *Pathophysiology* 2011; 18: 159-164.
 39. Sonawane RD, Vishwakarma SL, Lakshmi S, Rajani M, Padh H, Goyal RK. Amelioration of STZ-induced type 1 diabetic nephropathy by aqueous extract of *Enicostemma littorale* Blume and swertiamarin in rats. *Mol Cell Biochem* 2010; 340: 1-6.
 40. Sundaram R, Naresh R, Shanthi P, Sachdanandam P. Antihyperglycemic effect of iridoid glucoside, isolated from the leaves of *Vitex negundo* in streptozotocin-induced diabetic rats with special reference to glycoprotein components. *Phytomedicine* 2012; 19(3-4): 211-216.
 41. Sunil C, Ignacimuthu S, Agastian P. Antidiabetic effect of *Symplocos cochinchinensis* (Lour.) S. Moore. in type 2 diabetic rats. *J Ethnopharmacol* 2011; 134: 298-304.
 42. Thirumalai T, Therasa VS, Elumalai EK, David E. Hypoglycemic effect of *Brassica juncea* (seeds) on streptozotocin induced diabetic male albino rat. *Asian Pac J Trop Biomed* 2011; 4: 323-325.

43. Wadkar KA, Magdum CS, Patil SS, Naikwade NS. Antidiabetic potential and Indian medicinal plants. *J Herbal Med Toxicol* 2008; 2: 45-50.
44. Xin-Zhong Hu, Xiao-Hui Xing, Zheng-Mao Zhang, Rui-Qin Wu, Qingbin Guo,

Steve W. Cui, et al. Antioxidant effects of *Artemis sphaerocephala* Krasch. gum, on streptozotocin- induced type 2 diabetic rats. *Food Hydrocolloids* 2011; 25: 207-213.